

Sutab Colonoscopy Prep Instructions

What you need:

1. Sutab prep kit. Prescription. **Do not follow the instructions on the Sutab box.**
2. Two Simethicone anti-gas tablets. (Gas X or Phazyme) Purchase over the counter.

One (1) Day before your Procedure:

Step 1. Start a clear liquid diet beginning the day prior to your scheduled procedure. Examples include water, flavored water, sports drinks, sodas, fruit juices without pulp, Popsicles, Jell-O, clear broth, tea without cream or milk, and coffee without cream or milk.

**Avoid liquids that are red, orange or purple in color.*

**Remember to avoid solid food and non-clear liquids including Boost and similar protein drinks.*

**Please continue to drink the entire prep even if your stool looks clear or you feel cleaned out.*

Step 2. At 6:00 pm, take one simethicone gas tablet, open one bottle of 12 tablets. Fill the provided container with 16 ounces of water (up to the fill line). Swallow 1 tablet every 1 to 2 minutes. You should finish the 12 tablets and the entire 16 ounces of water within 20 minutes.

Important: If you experience preparation-related symptoms (examples: nausea, bloating, or cramping), pause or slow the rate of drinking the additional water until your symptoms diminish.

Step 3. Approximately one hour after the last tablet is ingested, fill the provided container a second time with 16 ounces of water (up to the fill line) and drink the entire amount over 30 minutes.

Step 4. Approximately 30 minutes after finishing the second container of water, fill the provided container again with 16 ounces of water (up to the fill line) and drink the entire amount over 30 minutes.

Step 5. Continue to hydrate with clear liquids before bedtime.

Step 5. Get some sleep.

Step 6. At (5 hours before arrival time), open the second bottle of 12 tablets. Fill the provided container with 16 ounces of water (up to the fill line). Swallow 1 tablet every 1 to 2 minutes. You should finish the 12 tablets and the entire 16 ounces of water within 20 minutes.

Important: If you experience preparation-related symptoms (for examples, nausea, bloating, or cramping), pause or slow the rate of drinking the additional water until your symptoms diminish.

Step 7. Approximately one hour after the last tablet is ingested, fill the provided container a second time with 16 ounces of water (up to the fill line) and drink the entire amount over 30 minutes.

Step 8. Approximately 30 minutes after finishing the second container of water, fill the provided container again with 16 ounces of water (up to the fill line) and drink the entire amount over 30 minutes. *This is a good time to take your usual morning meds. Delay taking insulin and oral diabetes meds until after your procedure.*

Your bowel prep is now complete! Do not take anything else by mouth until after your procedure.

Please call BGA at 205-271-8000 between 8:30am and 5:00pm **at least 48 hours before your procedure** if you have any questions or need to reschedule.

Colonoscopy Instruction Sheet

Purpose: This is an endoscopic evaluation of your large intestine (colon) and rectum to screen or recheck for polyps and colon cancer and/or investigate abnormal gastrointestinal symptoms.

Planning: Make sure you have your colonoscopy bowel preparation **at least 48 hours before your procedure**. If you do not, call BGA at 205-271-8000 during regular business hours.

- Plan to be off work for the day of your procedure.
- Arrange for your driver to take you to your procedure, wait for you while you are there and then drive you home after the procedure. You should not drive on the day of your procedure.
- Do not take your diabetes medications on the morning of your procedure.
- It is OK to continue taking aspirin, Tylenol, and anti-inflammatories.
- Stop taking iron supplements 3 days before your procedure.
- Start a low-residue diet beginning 3 days before your procedure. Avoid tough meats, whole grains, seeds, nuts, popcorn, raw vegetables, vegetable skins, fruit with seeds or skins, broccoli, cabbage, lettuce, corn, beans, and peas.
- Plan to abstain from chewing gum, mints, or tobacco products of any kind after midnight before your procedure.
- For your safety, please remove all metal including piercings and jewelry prior to your arrival.
- If you take a blood thinner for any reason and were not instructed to hold it in advance of your colonoscopy, please notify our office.

Blood Thinners:

-Coumadin (Warfarin/Jantoven)- 5 days prior
-Aggrenox (Aspirin-Dipyridamole) 5-7 days prior
- Brilinta (Ticagrelor)- 5 days prior
-Xarelto (Rivaroxaban)- 24 hours prior
-Lovenox (Enoxaparin)-24 hours prior
-Bevyxxa (Betrixaban)-48 hours prior
-Ticlid (Ticlopidine)-7 days prior

-Plavix (Clopidogrel)- 7 days prior
-Effient (Prasugrel)- 7 days prior
-Pradaxa (Dabigatran)- 24 hours prior
-Eliquis (Apixaban)-48 hours prior
-Pletal (Cilostazol)- 48 hours prior
-Savaysa (Edoxaban)-48 hours prior
-Zontivity (Vorapaxar)-5-7 days prior

Diabetes and Weight Management Medications:

Victoza (Liraglutide)- 24 hours prior
Saxenda (Liraglutide)- 24 hours prior
Xultophy (Liraglutide with Insulin)-24 hours prior
Gattex (Teduglutide)- 24 hours prior
Rybelsus (Semaglutide)- 24 hours prior
Byetta (Exenatide)- 24 hours prior

Bydureon BCise (Exenatide)- 7 days prior
Ozempic (Semaglutide)- 7 days prior
Wegovy (Semaglutide)-7 days prior
Trulicity (Dulaglutide)-7 days prior
Mounjaro (Tirzepatide)- 7 days prior

Prep: This is the process of ingesting clear liquids along with a strong laxative bowel-cleansing solution leading up to your colonoscopy. The laxative bowel prep is taken as a “split-dose.” This means that you take the first dose the night before your procedure and the second dose on the day of your procedure. This is very important to clear the colon of all waste material so that its surfaces can be carefully inspected. By the conclusion of your prep, your bowel movements will resemble clear to yellow water. **This step is extremely important! A suboptimal prep may result in your procedure being rescheduled, more frequent procedures or even missed lesions!**

Please call BGA at 205-271-8000 between 8:30am and 5:00pm **at least 48 hours before your procedure** if you have any questions or need to reschedule.